

TANAGRA MELANICTERA.

CHARACTER GENERICUS.

*Rostrum conicum, acuminatum, emarginatum,
basi subtrigonum, apice declive.*

Lin. Syst. Nat. p. 313.

CHARACTER SPECIFICUS, &c.

TANAGRA ferrugineo-fusca subtus lutea, pi-
leo genisque nigris, alis albo longitudina-
liter striatis.

Lath. Ind. Orn. p. 423.

TANAGRA supra ferruginea, subtus flavissi-
ma, capite nuchaque atris, alis albido-
striatis caudaque fuscis.

Lin. Syst. Nat. Gmel. p. 898.

TANAGRA melaniætera.

Guldenst. Nov. Comm. Petrop. 19. p. 465.

A Domino Gueldenstaedt in Actis Petropolitanis
ita describitur Tanagra melaniætera.

“ Magnitudine Tanagra melaniætera Emberizæ mi-
liariæ æqualis est. Caput supra a fronte usque ad
nucham et ad latera atrum ; collum supra et dorsum
brunneo-ferrugineum ; uropygium lutescenti-ferrugi-
neum ; avis tota subtus uniformiter flavissima. Alæ
complicatæ caudæ medium attingentes, fuscæ, albido
longitudinaliter striatæ ; remiges fuscæ, marginibus
albidis ; tectrices superiores remigibus concolores, in-

feriores autem albido-flavæ. Cauda subforcipata, tres pollices longa; rectrices duodecim, fuscæ, marginibus albido-flavicantibus.

“ Femina differt coloribus capitis et trunci; illa nimirum supra tota a fronte ad caudam usque sordide olivaceo-ferruginea fusco maculata, subtus tota ex albo flava.

“ Habitat in submontanis promontorii utriusque, et septentrionalis et meridionalis Caucasi, circa thermas ad flumen Terek obvias et in Georgia circa Teslium.”

563



Oriolus chinensis
Gmelin

BLACK-CROWNED TANAGER.

GENERIC CHARACTER.

Bill conic, acuminated, a little inclining towards the point; the upper mandible slightly ridged, and notched near the end.

SPECIFIC CHARACTER, &c.

Ferruginous-brown TANAGER, yellow beneath, with black crown and cheeks, and wings streaked with white.

Black-crowned TANAGER.

Lath. Syn. 2. p. 223.

This species is described by Guldenstadt in the Petersburgh Transactions. Its size is that of the common Bunting: the crown and cheeks are black, the neck and back brown-ferruginous, the rump yellowish-ferruginous, and the whole bird beneath deep yellow; the wings which, when closed, reach to the middle of the tail, are streaked longitudinally with white; the tail is slightly forked, measuring about three inches in length. The female differs in being entirely of a dull olive-ferruginous above, spotted with black, and beneath yellowish white.

It is an inhabitant of the Caucasian mountains and of the country of Georgia.